

***Review of Literature Relating to SaPII Formation and Capsid Size***  
**Meredith Hickson**

- Alouf J.E., and Muller-Alouf H. (2003) Staphylococcal and streptococcal superantigens: Molecular, biological and clinical aspects. *Int J Med Microbiology* **292**: 429-440.
- CDC. (2008) HIV/AIDS surveillance report, 2006. vol. 18.
- Charpentier E., Anton A.I., Barry P., Alfonso B., Fang Y., and Novick R.P. (2004) Novel Cassette-Based Shuttle Vector System for Gram-Positive Bacteria. *App Environ Microbiol* **70**: 6076-6085.
- Chen J., and Novick R.P. (2009) Phage-mediated intergeneric transfer of toxin genes. *Science* **323**: 139-141.
- Drake J.W. (1966) Ultraviolet Mutagenesis in Bacteriophage T4. *American Society for Microbiology* **91**: 1775-1780.
- Kelvins R.M., Morrison M.A., Nadle J., Petit S., Gershman K., Ray S., *et al.* (2007) Invasive methicillin-resistant *Staphylococcus aureus* infections in the united states. *JAMA* **298**: 1763-1771.
- King J., and Chiu W. (1997) The procapsid-to-capsid transition in double-stranded DNA bacteriophages. In *Structural Biology of Viruses*. Chiu W., Burnett R.M., and Garcea R.L. (eds). New York: Oxford University Press, pp. 288-311.
- Lindsay J.A., Ruzin A., Ross H.F., Kurepina N., and Novick R.P. (1998) The gene for toxin shock toxin is carried by a family of mobile pathogenicity islands in *Staphylococcus aureus*. *Mol Microbiol* **29**: 527-543.
- Maiques E., Ubeda C., Tormo M.A., Ferrer M.D., Lasa I., Novick R.P., and Penades J.R. (2007) Role of staphylococcal phage and SaPI integrase in intra- and interspecies SaPI transfer. *J Bacteriol* **189**: 5608-5616.
- Maiques E., Ubeda C., Campoy S., Salvador N., Lasa I., Novick R.R., *et al.* (2006) Betalactam antibiotics induce the SOS response and horizontal transfer of virulence factors in *Staphylococcus aureus*. *J Bacteriol* **188**: 2726-2729.
- Novick R.P., and Subedi A. (2007) The SaPIs: Mobile pathogenicity islands of *Staphylococcus*. *Chem Innumol Allergy* **93**: 42-57.
- Poliakov A., Chang J.R., Spilman M.S., Damle P.K., Christie G.E., Mobley J., and Dokland T. (2008) Capsid size determination by *Staphylococcus aureus* pathogenicity island SaPII involves specific incorporation of SaPII proteins into

- procapsids. *Journal of Molecular Biology* **380**: 465-475.
- Ruzin A., Lindsay J., and Novick R.P. (2001) Molecular genetics of SaPI1—a mobile pathogenicity island in *Staphylococcus aureus*. *Mol Microbiol* **41**: 365-377.
- Tropp B.E. 2008. *Molecular Biology: Genes to Proteins*. Boston (MA): Jones and Barlett. p. 290-298.
- Ubeda C., Maiques E., Knecht E., Lasa I., Novick R.P., and Pendades J.R. (2005) Antibiotic-induced SOS response promotes horizontal dissemination of pathogenicity island-encoded virulence factors in staphylococci. *Mol Microbiol* **56**: 836-844.
- Ubeda C., Maiques E., Tormo M., Campoy S., Lasa I., Barbe J., *et al.* (2007) SaPI operon I is required for SaPI packaging and is controlled by LexA. *Mol Microbiol* **65**: 41-50.